

## Conflict of interest

The authors declared that they do not have anything to disclose regarding funding or conflict of interest with respect to this manuscript.

## References

- [1] Romeo S, Kozlitina J, Xing C, Pertsemlidis A, Cox D, Pennacchio LA, et al. Genetic variation in PNPLA3 confers susceptibility to nonalcoholic fatty liver disease. *Nat Genet* 2008;40:1461–1465.
- [2] Valenti L, Al-Serri A, Daly AK, Galmuzzi E, Rametta R, Dongiovanni P, et al. Homozygosity for the PNPLA3/adiponutrin I148M polymorphism influences liver fibrosis in patients with nonalcoholic fatty liver disease. *Hepatology* 2010;51:1209–1217.
- [3] Valenti L, Alisi A, Galmuzzi E, Bartoli A, Del Menico B, Alterio A, et al. I148M patatin-like phospholipase domain-containing 3 gene variant and severity of pediatric nonalcoholic fatty liver disease. *Hepatology* 2010;52:1274–1280.
- [4] Tian C, Stokowski RP, Kershenovich D, Ballinger DG, Hinds DA. Variant in PNPLA3 is associated with alcoholic liver disease. *Nat Genet* 2009;42:21–23.
- [5] Stickel F, Buch S, Lau K, Zu Schwabedissen HM, Berg T, Ridinger M, et al. Genetic variation in the PNPLA3 gene is associated with alcoholic liver injury in caucasians. *Hepatology* 2011;53:86–95.
- [6] Cai T, Dufour JF, Muellhaupt B, Gerlach T, Heim M, Moradpour D, et al. Viral genotype-specific role of PNPLA3, PPARG, MTP and IL28B in hepatitis C virus-associated steatosis. *J Hepatol* 2011;55:529–535.
- [7] Valenti L, Rumi M, Galmuzzi E, Aghemo A, Del Menico B, De Nicola S, et al. Patatin-Like phospholipase domain-containing 3 I148M polymorphism, steatosis, and liver damage in chronic hepatitis C. *Hepatology* 2011;53:791–799.
- [8] Valenti L, Colombo M, Fargion S. PNPLA3 genotype and hepatocellular carcinoma in chronic hepatitis C. *Hepatology* 2011;53:1777.

- [9] Muller T, Buch S, Berg T, Hampe J, Stickel F. Distinct, alcohol-modulated effects of PNPLA3 genotype on progression of chronic hepatitis C. *J Hepatol* 2011;55:732–733.
- [10] Valenti L, De Feo T, Fracanzani AL, Fatta E, Salvagnini M, Arico S, et al. Cytotoxic T-lymphocyte antigen-4 A49G polymorphism is associated with susceptibility to and severity of alcoholic liver disease in Italian patients. *Alcohol Alcohol* 2004;39:276–280.

Luca Valenti\*

Università degli Studi di Milano, Fondazione IRCCS Ca' Granda,  
Ospedale Maggiore Policlinico, Pad. Granelli, via F. Sforza 35, 20122  
Milan, Italy  
Tel.: +39 02 503 20278; fax: +39 02 503 20296  
\*E-mail address: luca.valenti@unimi.itm

Massimo Colombo

Department of Internal Medicine,  
A.M. Migliavacca Center for Liver Disease,  
First Division of Gastroenterology, Università degli Studi di Milano,  
Fondazione IRCCS "Ca' Granda" Ospedale Maggiore Policlinico,  
Milan, Italy

Silvia Fargion

Department of Internal Medicine,  
A.M. Migliavacca Center for Liver Disease,  
First Division of Gastroenterology, Università degli Studi di Milano,  
Fondazione IRCCS "Ca' Granda" Ospedale Maggiore Policlinico,  
Milan, Italy

## Reply to: “Modulation of the effect of PNPLA3 I148M mutation on steatosis and liver damage by alcohol intake in patients with chronic hepatitis C”

**PNPLA3 rs738409 and fibrosis progression in chronic hepatitis C – There is more to it than just fat!**

To the Editor:

The Hepatology community has seen a substantial rise in publications reporting data on genetic risk factors conferring risk to fibrosis progression in non-alcoholic, alcoholic, and chronic hepatitis C-related liver diseases. For long, hypothesis-driven, often single-centre genetic case control studies have reported associations of certain candidate genes with fibrosis/cirrhosis which could not be replicated. With genome-wide association studies (GWAS) becoming a standard tool in translational research, novel candidate genes enter the stage which have been identified by systematic screening experiments [1]. GWAS make use of a hypothesis-free, or rather – hypothesis-generating approach, and have the potential to uncover genetic risk factors which had not been considered based on our previous pathophysiological understanding. A striking example for such revelation is the gene coding for patatin-like phospholipase domain containing-3 (PNPLA3; adiponutrin) of which a polymorphic variant (rs738409 G > C) was found associated with liver fat content in a landmark GWAS [2], and subsequently with progression of non-alcoholic fatty liver disease (NAFLD) [3], and alcoholic liver disease (ALD) [4,5] by means of

candidate case control studies. The role of PNPLA3 rs738409 as a risk factor for progressive fibrosis can now be extended to chronic hepatitis C. Only in 2011, four studies presented data on the role of PNPLA3 rs738409 G which unanimously confirm that carriage of at least one allele increases the risk of advanced fibrosis and cirrhosis [6–9] (Table 1). Our own data indicate that this association is particularly evident in hepatitis C virus-infected individuals who regularly drink alcohol as opposed to abstainers [6]. Regarding the latter, PNPLA3 rs738409 G appears to modulate fat storage in the liver by a yet unknown mechanism, possibly by interaction of PNPLA3 with viral epitopes. Indeed, from the data presented in published studies, it becomes clear that PNPLA3 rs738409 G neither aggravates steatosis nor fibrosis in patients with genotype 3, supporting the hypothesis that mostly genotype 1-specific interactions with PNPLA3 are involved.

In this issue of the *Journal*, Valenti and coworkers have undertaken a pooled analysis of our [6] and their data [7] after stratifying their patients into “at-risk” drinkers (daily alcohol consumption  $\geq 20$  g) and abstainers [10]. Hereby, they confirm that carriage of PNPLA3 rs738409 G is a risk factor for steatosis in abstainers, but not in at risk drinkers, possibly due to a confounding effect on steatosis by concomitant alcohol consumption. In addition, by increasing the numbers of total cases and controls,

## Letters to the Editor

**Table 1. Genetic candidate gene association studies on the role of *PNPLA3* rs738409 G as a risk factor for fibrosis and cirrhosis.**

Author [Ref]	Patients (n)	End points	Subgroup analysis		Results		
					OR	95% CI	p value
Müller <i>et al.</i> , [6]	Steatosis (442) Cirrhosis (605)	Steatosis Cirrhosis	Abstainers "At-risk" drinkers	Association with			
				• steatosis in abstainers • cirrhosis in "at-risk" drinkers	12.61 4.77	1.48-107.7 1.39-6.38	0.021 0.013
Valenti <i>et al.</i> , [7]	Two independent cohorts (325/494)	Steatosis	None	Association with			
		Cirrhosis		• steatosis	1.9	1.4-2.7	<0.001
		HCC		• fibrosis/cirrhosis	1.47	1.2-19	0.002
		Therapy response		• therapy response • HCC	0.63 2.61	0.4-0.8 1.3-3.6	0.006 0.002
Cai <i>et al.</i> , [8]	626	Steatosis	Genotype 3 Genotype non-3	Association with • steatosis but only in viral genotype non-3	1.88	1.57-2.25	<0.001
Trepo <i>et al.</i> , [9]	537	Steatosis	None	Association with			
		Fibrosis		• steatosis	2.55	1.08-6.03	0.034
		Fibrosis progression		• fibrosis	3.13	1.50-6.51	0.002
		Therapy response		• fibrosis progression	2.64	1.22-5.67	0.013

they convincingly show that *PNPLA3* rs738409 G is also a risk factor for cirrhosis in abstainers, with an additional increase of the odds ratio (OR) in at risk drinkers. Of note, a subgroup analysis of our latest data [9], confirmed these observations [6,10]. Again, only abstainers were at risk of steatosis (OR = 4.52 95% confidence interval [CI] 1.32–15.48,  $p = 0.016$ ). Moreover, *PNPLA3* rs738409 G remained associated with advanced fibrosis in abstainers (OR = 2.68 95% CI 1.18–6.06,  $p = 0.018$ ), but still its impact on fibrosis was stronger in "at-risk" drinkers (OR = 10.34 95% CI 1.15–93.16,  $p = 0.037$ ). In our view, these analyses provide a perfect example of host-environment interactions in which the risk of developing cirrhosis increases in a "dose-dependent" manner with the number of risk factors present (*PNPLA3* rs738409 G, hepatitis C virus, and alcohol).

Together, data from these four independent studies deliver strong evidence for a significant role of *PNPLA3* rs738409 for progressive liver disease not only in NAFLD and ALD, but also in CHC. What needs to be elucidated now is to decipher the cross-talk between *PNPLA3* (and its genetic variation) and fibrosis-related intracellular signalling. Fat accumulation may do harm in hepatitis C infected subjects – but as we know from NAFLD, it requires a "second hit" to cause inflammation and fibrosis.

### Conflict of interest

The authors declared that they do not have anything to disclose regarding funding or conflict of interest with respect to this manuscript.

### References

- [1] Zimmer V, Lammert F. Genetics in liver disease: new concepts. *Curr Opin Gastroenterol* 2011;27:231–239.
- [2] Romeo S, Kozlitina J, Xing C, Pertsemlidis A, Cox D, Pennacchio LA, et al. Genetic variation in *PNPLA3* confers susceptibility to non-alcoholic fatty liver disease. *Nat Genet* 2008;40:1461–1465.
- [3] Valenti L, Al-Serri A, Daly AK, Galmozzi E, Rametta R, Dongiovanni P, et al. Homozygosity for the *PNPLA3*/adiponutrin I148M polymorphism influences liver fibrosis in patients with nonalcoholic fatty liver disease. *Hepatology* 2010;51:1209–1217.
- [4] Stickel F, Buch S, Lau K, Zu Schwabedissen HM, Berg T, Ridinger M, et al. Genetic variation in the *PNPLA3* gene is associated with alcoholic liver injury in Caucasians. *Hepatology* 2011;53:86–95.
- [5] Trépo E, Gustot T, Degre D, Lemmers A, Verset L, Demetter P, et al. Common polymorphism in the *PNPLA3*/adiponutrin gene confers higher risk of cirrhosis and liver damage in alcoholic liver disease. *J Hepatol* 2011;54:906–912.
- [6] Müller T, Buch S, Berg T, Hampe J, Stickel F. Distinct, alcohol-modulated effects of *PNPLA3* genotype on progression of chronic hepatitis C. *J Hepatol* 2011;55:729–733.
- [7] Valenti L, Rumi MG, Galmozzi E, Aghemo A, Del Menico B, De Nicola S, et al. *PNPLA3* I148M polymorphism, steatosis, and liver damage in chronic hepatitis C. *Hepatology* 2011;53:791–799.
- [8] Cai T, Dufour JF, Muellhaupt B, Gerlach T, Heim M, Moradpour D, et al. Viral genotype-specific role of *PNPLA3*, *PPARG*, *MTTP* and *IL28B* in hepatitis C virus-associated steatosis. *J Hepatol* 2011;55:529–535.
- [9] Trépo E, Pradat P, Potthoff A, Momozawa Y, Quertinmont E, Gustot T, et al. Impact of *PNPLA3* (rs738409 C>G) polymorphism on fibrosis progression and steatosis in chronic hepatitis C. *Hepatology* 2011;54:60–69.
- [10] Valenti L, Colombo M, Fargion S. Modulation of the effect of *PNPLA3* I148M mutation on steatosis and liver damage by alcohol intake in patients with chronic hepatitis C. *J Hepatol* 2011;55:1471–1472.

Felix Stickel\*

Department of Internal Medicine I,  
University Hospital Schleswig-Holstein, Kiel,  
Germany

Department of Visceral Surgery and Medicine,  
Institute of Clinical Pharmacology and Visceral Research,  
Inselspital, University of Bern, Murtenstr. 35, 3010 Bern,  
Switzerland

\*E-mail address: felix.stickel@ikp.unibe.ch

Eric Trépo

Christophe Moreno

Department of Gastroenterology,  
Hepatopancreatology and Digestive Oncology, Erasme Hospital,  
Université Libre de Bruxelles, Brussels, Belgium  
Laboratory of Experimental Gastroenterology,  
Université Libre de Bruxelles, Brussels, Belgium

Jochen Hampe

Department of Internal Medicine I,  
University Hospital Schleswig-Holstein, Kiel,  
Germany